

INVARIANTS IN $\mathcal{C} \cup \mathcal{P}$ -GEOMETRY

GUY BURDET

ABSTRACT. This note describes the construction of $\mathcal{C} \cup \mathcal{P}$ -invariant differential operators on statistical manifolds, i.e. of operators canonically associated to a geometry which synthetizes the properties of conformal and projective geometries.

1. INTRODUCTION

Let (S, g, t) be a statistical manifold, i.e. a Riemannian manifold (S, g) endowed with a fully symmetric 3 covariant tensor field t , the *skewness tensor*. This tensor has been introduced to formalize the notion of statistical curvature through the introduction of a family $\{\overset{\alpha}{\nabla}\}_{\alpha \in \mathbb{R}}$ of affine connections defined as follows [1][2]

$$\overset{\alpha}{\nabla}_x Y = \overset{o}{\nabla}_x Y - \frac{\alpha}{2}(t \cdot g^{-1})(X, Y), \quad (1.1)$$

for any couple X, Y of vector fields over S , $\overset{o}{\nabla}$ denoting the canonical Levi-Civita connection and the dot $(\cdot)$ an obvious contraction. In fact $\overset{\alpha}{\nabla}$ is the unique torsion-free connection satisfying

$$\overset{\alpha}{\nabla} g = \alpha t, \quad (1.2)$$

and $(\overset{\alpha}{\nabla}, g)$ form a Codazzi pair [3], i.e. Codazzi equations are satisfied

$$(\overset{\alpha}{\nabla}_x g)(Y, Z) = (\overset{\alpha}{\nabla}_y g)(X, Z).$$

In a previous paper [4], we have dealt with the so-called $\mathcal{C} \cup \mathcal{P}$ -geometry, where the $\mathcal{C} \cup \mathcal{P}$ symbol refers to a conformal implementation of the projective geometry in which the properties of both conformal and projective geometries are preserved, as it can be seen from the following

Definition. Let α be a real number. Statistical manifolds (S, g, t) and $(S, \tilde{g}, \tilde{t})$ are α - $\mathcal{C} \cup \mathcal{P}$ -related if, for some positive smooth function η over S ,

$$\tilde{g} = \eta g, \quad \tilde{t} = \eta(t + \text{sym}(g \otimes \psi))$$

where ψ is an exact 1-form satisfying $d(\text{Log } \eta) = -\alpha \psi$.

and

1991 *Mathematics Subject Classification.* 53 C 99.

Proposition. *For any real α , the families of α -connections of two α -cUp-related statistical manifolds are projectively related, i.e. the corresponding 1-form connections satisfy the following relation*

$$\tilde{\varphi}^\alpha - \varphi^\alpha = -\alpha (\theta \otimes \psi + (\psi \cdot \theta) Id), \quad (1.3)$$

and the curvature 2-forms are such that

$$\tilde{\Phi}^\alpha - \Phi^\alpha = \alpha \theta \wedge \overset{\alpha}{\nabla} \psi + \alpha^2 (\theta \otimes \psi) \wedge (\psi \cdot \theta). \quad (1.4)$$

The invariants of the cUp-structure themselves are described in [4], here we want to study the closely related question of invariance of auxilliary objects associated to the cUp-structure.

2. CUP-INVARIANT SECOND ORDER DIFFERENTIAL OPERATORS

As the conformal one's, the cUp-geometry depends on the (conformal) η factor only, so we take up the step traced by the conformal experience by saying that a differential operator D , depending on a choice of the couple (g, t) , is *cUp-invariant of type $(r; s)$* if the operator $\tilde{D}$, corresponding to the rescaled metric $\tilde{g}$ and to the modified tensor $\tilde{t}$, is such that $\tilde{D}\tilde{f} = \eta^s Df$, where f is a *density of weight r* , section of a suitable line bundle, and $\tilde{f} = \eta^r f$, $r, s \in \mathbb{R}$.

Let us consider the α -Hessian $\overset{\alpha}{\nabla} \otimes d$ acting on functions on S , we look for the existence of a modification of it, form invariant for the cUp-geometry. Knowing that conformal invariant operators may be expressed in terms of the Levi-Civita connection and the Ricci curvature of a metric in the conformal class [5], we study the operator $\overset{\alpha}{\nabla} \otimes d + k Ric^\alpha$, $k \in \mathbb{R}$, where Ric^α denotes the Ricci tensor of the α -connection.

Starting from Rel(1.4) it can be shown that the Ricci tensors of two α -cUp-related connections are such that

$$\tilde{Ric}^\alpha = Ric^\alpha + (n-1)\alpha\{\overset{\alpha}{\nabla} \psi + \alpha(\psi \otimes \psi)\}, \quad (2.1)$$

where $n = \dim S$.

Then, by using Rel(1.3) one gets

$$(\tilde{\nabla}^\alpha \otimes d)(\eta^r f) = \eta^r (\nabla^\alpha \otimes d)f + f(\nabla^\alpha \otimes d)\eta^r + d(\eta^r) \otimes \nabla^\alpha f + df \otimes \nabla^\alpha(\eta^r), \quad (2.2)$$

taking into account for $d(\text{Log} \eta) = -\alpha \psi$, one establishes the following

Proposition. *The modified Hessian $\overset{\alpha}{\nabla} \otimes d + k Ric^\alpha$ is α -cUp-invariant of type $(1; 1)$ for $k = \frac{1}{n-1}$:*

$$(\tilde{\nabla}^\alpha \otimes d + \frac{1}{n-1} \tilde{Ric}^\alpha)\tilde{f} = \eta(\overset{\alpha}{\nabla} \otimes d + \frac{1}{n-1} Ric^\alpha)f. \quad (2.3)$$

Remark The integrability conditions of the differential system $(\overset{\alpha}{\nabla} \otimes d + k Ric^\alpha)f = 0$ are given by

$$Riem^\alpha = k(Id \otimes Ric^\alpha - Ric^\alpha \otimes Id).$$

Except for the flat case $Riem^\alpha = 0$, this property is verified to hold, by taking the trace, for the cUp-invariant Hessian only, and for statistical manifolds such that

$$Riem^\alpha = \frac{1}{n-1}(Id \otimes Ric^\alpha - Ric^\alpha \otimes Id). \quad (2.4)$$

For instance this property is satisfied for the statistical manifold associated to the gaussian law and for the statistical manifold of multinomial distributions.

3. THE CUP-LAPLACIAN

Let us consider the trace of the Ricci modified Hessian operator

$$g^{-1}(\overset{\alpha}{\nabla} \otimes d + k Ric^\alpha) = g^{-1}(\overset{\alpha}{\nabla} \otimes d) + k R^\alpha$$

where R^α denotes the *scalar α -curvature* $R^\alpha = g^{-1}(Ric^\alpha)$. This expression can be written by using the α -Laplacian $= \text{div}^\alpha \text{grad}$, denoted by $\overset{\alpha}{\Delta}$, but we have to take into account for the non-metricity of the connection; so we are led to give the

Definition. *The second order differential operator given by*

$$\Delta_{cUp} = g^{-1}(\overset{\alpha}{\nabla} \otimes d + \frac{1}{n-1} Ric^\alpha) = \overset{\alpha}{\Delta} + \alpha g^{-1}.t.g^{-1}.d + \frac{1}{n-1} R^\alpha, \quad (3.1)$$

where each point denotes an obvious contraction due to the symmetry of implied tensor fields, is called the *cUp - Laplacian*.

Indeed this operator is associated to a cUp -structure canonically as it is clear from the following

Proposition.

$$\tilde{\Delta}_{cUp} \tilde{f} = \Delta_{cUp} f, \quad (\tilde{f} = \eta f), \quad (3.2)$$

the *cUp - Laplacian* is a *cUp - invariant differential operator of type (1;0)*.

This property stems from Rel(2.3) by noting that $\tilde{g}^{-1} = \frac{1}{\eta}g^{-1}$.

4. QUASI-CUP-INVARIANT NON-LINEAR OPERATORS

We now wish to construct a non-linearly modified cUp- Laplacien through a self-interaction term f^a with a coupling potential λ , function on S . We can use property (3.2) to get immediately the

Proposition.

$$\tilde{\Delta}_{cUp} \tilde{f} + \tilde{\lambda} \tilde{f}^a = \Delta_{cUp} f + \lambda f^a, \quad (\tilde{f} = \eta f, \tilde{\lambda} = \lambda/\eta^a, a = \text{constante} \neq 0). \quad (4.1)$$

It is worthnoticing that there is no constraint over the non-linearity coefficient a contrarily to the conformal case for which a depends on the dimension of the manifold [6].

REFERENCES

1. N.N. Chentsov, *Statistical Decision and Optimal Inference*, “Nauka”, Moscow, 1972 (Russian); English transl., American Mathematical Society, vol. 53, Providence, R. I., 1982..
2. S-i.Amari, *Differential Geometrical Methods in Statistics*, Lecture Notes in Statistics., vol. 28, Springer-Verlag, New York, 1985.
3. U. Pinkall, A. Schwenk-Schellschmidt, U. Simon, *Geometric methods for solving Codazzi and Monge-Ampere equations*, Mathematische Annalen, **298** (1994), 89–100.
4. G. Burdet, M. Perrin,, *Conformal Implementation of the Projective Equivalence over Statistical Manifolds*, Indagationes Mathematicae, **8** (1997), 7–17.
5. R.J. Baston, *Verma modules and differential conformal invariants*, J. Differential Geometry, **32** (1990), 851–898.
6. B. Ørsted, *Conformally Invariant Differential Equations and Projective Geometry*, Journal of Functional Analysis **44** (1981), 1–23.

CENTRE DE PHYSIQUE THÉORIQUE, CNRS-LUMINY, CASE 907, F-13288 MARSEILLE, FRANCE
E-mail address: `burdet@cptsu2.univ-mrs.fr`